

A retrospective study on frequency of canine tumors at the Veterinary Teaching Hospital in Rabat (Morocco)

Soraya NOURY^{1,2}, Hassan BOUAYAD², Noursaid TLIGUI¹, Rahma AZRIB²

Abstract

¹ Department of Veterinary Pathology and Public Health, IAV Hassan II, Rabat, Morocco

² Veterinary Teaching Hospital, IAV Hassan II, Rabat, Morocco

* Corresponding author

sora.noury@gmail.com

Received 07/09/2019

Accepted 16/12/2019

This work is a contribution to the study of tumors in dogs at the Veterinary Teaching Hospital of the Hassan II Agronomic and Veterinary Medicine Institute in Rabat, Morocco. The study aimed to present a summary of the relative frequency of tumors in dogs, their preferred localization and the influence of age, sex and breed. During the study period from 2004 to 2017, a total of 3099 records were consulted, among them 166 cases of tumors in dogs were diagnosed in both sexes, male and female, whose ages were between 3 months to 15 years. Tumors in dogs have a frequency of approximately 5.4% over the entire canine disease diagnosed at the Veterinary Teaching Hospital. The average age group of onset of these tumors are middle-aged dogs of 7-12 years (65.7%). The German shepherd (30.7%), Poodle (22.9%), Labrador (10.2%), Rottweiler (5.4%) and Pekinese (3.6%) would be more prone to tumors. The most common tumors originated from the mammary gland (46.4%) and the skin and subcutis (25.9%). To our knowledge, there was no study like this one considering canine tumors in Morocco. Therefore, this study provides an important preliminary step to characterize and understand the occurrence of canine tumors in Rabat region.

Keywords: Tumor, dog, frequency, sex, age, breed, Morocco

INTRODUCTION

Tumors in pets represent one of the main reasons for consultation and are among the most important causes of morbidity and mortality in the canine and feline species. This condition has been the subject of numerous investigations around the world. Several previous animal tumor registries are still active today (Šoštarić-Zuckermann *et al.*, 2013) and include Veterinary Medical Database (VMDB, 2010), VetCancer Registry, Norwegian Canine Cancer Registry (Gamlem *et al.*, 2008), Danish Veterinary Cancer Registry (Brønden *et al.*, 2010), Animal Tumor Registry of the Vicenza and Venice Provinces of the Veneto Region (Vascellari *et al.*, 2009) and Animal Tumor Registry of Genoa (Merlo *et al.*, 2008). One of the primary objectives of these registries is to obtain an incidence rate of tumors per year per 100,000 dogs. To obtain this figure, it is necessary to know the exact population of dogs in the study area (Šoštarić-Zuckermann *et al.*, 2013; Merlo *et al.*, 2008).

In Morocco, we do not yet have a comprehensive national cancer database that provides a perfect understanding of the incidence and characteristics of cancer in canine and feline species. Moreover, the veterinarian is often confronted with this pathology, but despite significant progress in management, many animals remain without diagnosis and therefore without treatment. Most bibliographic data on tumors in dogs come from studies that focus on clinical and histopathological description and concern animals being treated. In this work, preliminary information on the frequency of different tumor types in dogs at the veterinary teaching hospital in Rabat, Morocco will be analyzed.

The objective of this study is to describe the frequency of tumors in dogs but also the types of tumors diagnosed. The study also evaluates the influence of age, sex and breed to document factors likely to be the origin or incriminated in this pathology.

Compared with existing data, this study is the first of its kind which focuses on all tumor types in dogs, based on a population of dogs presented to consultations.

MATERIALS AND METHODS

Data source

In order to have an overview of the incidence of the malignant neoplasms in canine population in the region of Rabat, Morocco, a retrospective study was performed at the Veterinary Teaching Hospital of the Hassan II Institute of Agronomy and Veterinary Medicine (Rabat) during the period of 2004-2017. A total of 3099 handwritten documents, collected from the registers, were reviewed. It lists all the animals seen at the hospital and using samples from canine biopsy and necropsy samples that proved to be neoplastic and were sent to the Department of Histology and Pathological Anatomy. Diagnoses were reached by experienced veterinary pathologists performing histopathologic evaluation of standard Hematoxylin Eosine stained microscopic slides. The samples from surgeries within the hospital were taken from different anatomical sites (mammary, cutaneous...etc).

Data preparation

The datasets were compiled in an Excel file by year and organized according to anatomic location (site) of

the tumor and variables set for the investigation were recorded according to age, sex and breed of dog from which a tumor biopsy was taken, or from which the tumour was excised at the surgery or necropsy.

Statistical analyses

Statistical analyses of collected data were performed in Microsoft Office Excel in order to evaluate the distribution of tumors. All cases were sorted by breed, sex and three age groups of six years as follows: 0 -6 years, 7-12 years, 13 -15 years.

RESULTS

Among 3099 sick dogs presented to the hospital, a total of 166 tumors were diagnosed between 2004 and 2017, which represent a frequency of 5.4%. The most frequently diagnosed neoplasms were tumors of the mammary gland (46.4 %), skin and subcutis (25.9 %), oral cavity (6 %), and genital tract (6 %).

Sex

Of those tumors, 60 were in male dogs (36.1%) and 106 were in female dogs (63.8%). Among male dogs, the most frequently diagnosed neoplasms were tumors of the skin and subcutis (19.3%), genital system (4.8%) and oral cavity (3.6%). Among female dogs, tumors of the mammary gland (46.4%), skin and subcutis (6.6%), and liver (3%) were dominant. Detailed information considering the frequency of tumors in a particular body system and distribution in a given body system according to gender is presented in table 1.

Age

The studied population includes dogs whose age is between 3 months and 15 years. The incidence of tumors increases significantly in the middle aged dogs (65.7%), with a clear decrease beyond 12 years (10.8%). When considering distribution of tumors in various age groups, the incidence of tumors of the skin and

Table 1: Distribution of tumors by body system and gender

Body system	Male	Female	Total
Mammary gland	-	77 (46.4 %)	77 (46.4 %)
Skin and subcutis	32 (19.3 %)	11 (6.6 %)	42 (25.9 %)
Oral cavity	6 (3.6 %)	4 (2.4 %)	10 (6 %)
Genital tract	8 (4.8 %)	2 (1.2 %)	10 (6 %)
Liver	4 (2.4 %)	5 (3 %)	9 (5.4 %)
Bones and joints	2 (1.2 %)	4 (2.4 %)	6 (3.6 %)
Endocrine and exocrine glands	4 (2.4 %)	-	4 (2.4 %)
Hemopoietic cells and vascular system	3 (1.8 %)	1 (0.6 %)	4 (2.4 %)
Respiratory	-	1 (0.6 %)	1 (0.6 %)
Ocular	1 (0.6 %)	-	1 (0.6 %)
Gastric	-	1 (0.6 %)	1 (0.6 %)
Total	60 (36.1 %)	106 (63.8%)	166 (100%)




Figure 1: Distribution of tumors by body system and by age group




Figure 2: Distribution of tumors by body system and by dog breed

subcutis were highest in the youngest age groups, while in middle-aged and older (female) dogs, mammary tumors were clearly the most common tumors. Precise data concerning the latter is given in figure 1.

Breed

In the present study, 156 dogs affected by tumors were purebred (94% dogs), and 10 dogs were crossed (6% of dogs). The most common breed in this study was the German shepherd (30.7%), followed by Poodle (22.9%), Labrador (10.2%), Rottweiler (5.4%), and Pekinese (3.6%).

In all these breeds, the most common organ system affected with tumors was the mammary gland, with the exception of the Rottweiler and the Pekinese in which the most common tumors originated respectively from the skin and subcutis, and the bones and joints. The distributions of tumors in the most common breeds are shown in figure 2. Other breeds have not been compared because of their very small numbers.

DISCUSSION

The results of this study reveal not only interesting aspects but also limitations regarding canine tumor types and frequency. A total of 166 canine tumors reported during a twelve year period in the teaching veterinary hospital in Rabat is not a large sample when compared with the 1172 tumors reported during an eight-year study in northern of Italy (Baioni *et al.*, 2017) and 3378 tumors reported during the ten-year study in a veterinary teaching hospital in the region of Londrina, Brazil (Rossetto *et al.*, 2009). One reason for the relatively small number of samples was the cost of histopathological analysis. When looking at total numbers, our study represents just malignant tumors. This could be explained by the fact that most owners do not treat slow-growing neoplasm or small tumors of their pets.

The body systems mostly affected were mammary gland (46.4 %), skin and subcutis (25.9 %), followed by oral cavity (6 %) and genital tract (6 %). A similar proportion of tumors by body system has been reported by other authors (40.8% skin and subcutis, 34.8% mammary gland, and 7.2% genital (Vascellari *et al.*, 2009); 51% skin and subcutis, 30% mammary gland, and 3.9% genital (Gamlem *et al.*, 2008). It is widely accepted that the perceived frequencies of tumors in these body systems tend to be greater as they are easier to recognize by the owners and physical examination than are tumors of the respiratory tract, digestive tract, and other internal organs that require specific examinations such as X-ray, computed tomography scanning (CT), magnetic resonance imaging (MRI) or ultrasound examination (Merlo *et al.*, 2008). Smaller differences in the proportion of tumors by body system could be the result of different environmental influences or customs regarding spaying of animals between the studied populations and those from other studies.

The sex ratio in our study (36.1% male, 63.8% female dogs) is not equally distributed. This difference can be explained by the fact that mammary tumors that exclusively affect female dog, occupy a frequency of 46.4% of all neoplasms. Our result is supports many previous studies (Anderson and Johnson, 1967; Cohen *et al.*, 1974; Luttgen *et al.*, 1980; Hayes *et al.*, 1983; Bukowski *et al.*, 1998) which showed that females are more affected by neoplasms than males and this is due to the fact that mammary tumors are of great importance in female tumor pathology. According to our study, the males has a high incidence than females of tumors originated from skin and subcutis. This has been shown by a great number of authors (Cohen *et al.*, 1974; Hayes *et al.*, 1983; Bukowski *et al.*, 1998; Collas, 1997).

The studied population includes dogs whose age is between 3 months and 15 years. The incidence of tumors increased significantly in the middle aged dogs (65.7%), with a clear decrease beyond 12 years (10.8%). These values confirm previous work. Indeed, several studies (Cadiot, 1907; Dorn, 1967; Hayes, 1978; Prymak *et al.*, 1988; Mialot and Lagadic, 1990; Prange *et al.*, 1998) indicated that the incidence of tumors increases significantly with age until 9-10 years, remains constant around the age of 12 years and then decreases. Tumors are therefore diseases of the adult dog.

The distribution of tumors by age group shows skin and subcutis tumors predominate in the youngest age groups and mammary gland tumors are the most common tumors in middle-aged and older age groups in females. A similar distribution is presented by Gamlem *et al.*, (2008).

The distribution of tumors by breed showed breed predispositions. In our study, 156 dogs affected by tumors were purebred (94% dogs) and 10 dogs were crossed (6% of dogs). The most common breed affected in this study was the German shepherd (30.7%), followed by Poodle (22.9%), Labrador (10.2%), Rottweiler (5.4%), and Pekinese (3.6%). Previous studies have shown a predisposition due to racial polymorphism, although the dog populations initially studied differed according to authors. Thus, a previous study made by Priester (1979) shows that pure breeds are much more predisposed than crosses. Pet dog breeds such as Dachshund, Bichon, Poodle, Yorkshire and Pinscher are prone to mammary tumors (Mialot and Lagadic, 1990). Our study shows that both large breeds (German shepherd and Labrador) and pet dog breed (poodle) are prone to mammary tumors.

Large breeds are prone to skeletal tumors (Tjalma, 1966; Brodey, 1970). The high incidence of bones and joints in Pekinese was unexpected and should be the subject of further study.

Until now, there was no study similar to this one considering canine tumors in Morocco. Therefore, this study provides a primary step to better characterize and understand the occurrence of canine tumors in Rabat

region. Most of the incidences of tumors are comparable with those from other studies. However, the results and comparison with other studies indicate the need to conduct subsequent studies and the need for establishment of a computer based registry of tumors in domestic animals in Morocco.

ACKNOWLEDGEMENTS

The authors sincerely acknowledge Ms. ElBakour Najat of the LAV Hassan II Veterinary Teaching Hospital for her help in collecting the files needed for this study. Many thanks to Ms. Irene M. Struikma for her help in partially editing the text.

REFERENCES

- Anderson N.V., Johnson Kh. (1967). Pancreatic carcinoma in the dog. *J. Am. Vet. Med. Assoc.* 150: 286-295.
- Baioni E., Scanziani E., Vincenti M. C., Leschiera M., Bozzetta E., Pezzolato M., Desiato R., Bertolini S., Maurella C., Ru G. (2017). Estimating canine cancer incidence, findings from a population-based tumour registry in northwestern Italy. *BMC Veterinary Research* 13:203.
- Brodey R.S. (1970). Canine and feline neoplasia. *Adv Vet Sci Comp Med.* 14: 309-354.
- Brønden L. B., Nielsen S. S., Toft N., Kristensen A. T. (2010). Data from the Danish Veterinary Cancer Registry on the occurrence and distribution of neoplasms in dogs in Denmark. *Vet. Rec.* 166: 586-590.
- Bukowski J.A., Wartenberg D., Goldschmidt M. (1998). Environmental causes for sinonasal cancers in pet dogs and their usefulness as sentinels of indoor cancer risk. *J. Toxicol. Environ. Health.* 54 :579-591.
- Cadiot J. (1907). Sur le cancer des animaux. *Rec. Med. Vet.* 84: 5-25.
- Cohen D., Reif J.S., Brodey R.S., Keiser H. (1974). Epidemiological analysis of the most prevalent sites and types of canine neoplasia observed in a veterinary hospital. *Cancer Res.* 34: 2859-2868.
- Collas G. (1997). Chirurgie des cavités nasales et des sinus. In *Encyclopédie vétérinaire*, Elsevier, V. 2.
- Dorn Cr. (1967). The epidemiology of cancer in animals. *Calif. Med.* 107: 481-489.
- Gamlem H., Nordstoga K., Glatte E. (2008). Canine neoplasia - Introductory paper. *APMIS Suppl.* 125: 5-18.
- Hayes H.M. (1978). The comparative epidemiology of selected neoplasm between dogs, cats and humans. A review. *Europ. J. Cancer*, 14: 1299-1308.
- Hayes H.M., Morin M.M., Rubenstein D.A. (1983). Canine Biliary carcinoma: epidemiological comparisons with man. *J. Comp. Pathol.* 93: 99-107.
- Luttgen P.J., Braund K.G., Brawner Jr W.R., Vandeveld M. (1980). A retrospective study of 29 tumors in the dog and cat. *J. Small. Anim. Pract.* 21: 213-226.
- Merlo D. F., Rossi L., Pellegrino C., Ceppi M., Cardellino U., Capurro C., Ratto A., Sambusso P. L., Sestito V., Tanara G. (2008). Cancer incidence in pet dogs: findings of the animal tumor registry of Genoa, Italy. *J. Vet. Intern. Med.* 22: 976-984.
- Mialot M., Lagadic M. (1990). Epidémiologie descriptive des tumeurs du chien et chat. *Rec. Med. Vet.*, 166 : 937-947.
- Prange H., Falk-Junge G., Katenkamp D., Schneider E., Zieger M. (1998). Distribution, epizootiology and radiographical diagnosis of intrathoracic neoplasms in dogs. *Archiv. Exp. Veterinar Med.* 42: 637-649.
- Priester W.A. (1979). Epidemiology, in veterinary cancer medicine. *Lea And Febiger.* 1: 14-32.
- Prymak P., McKee L.J., Goldschmidt M.H., Glickman L.T. (1988). Epidemiological, Clinical, Pathologic, and prognostic characteristics of splenic hemangiosarcoma and splenic hematoma in dogs: 217 cases. *J. Am. Vet. Med. Assoc.* 193: 706-712.
- Rossetto V. J.V., Moreno K., Grotti C.B., A.C.F.D. Reis, Bracarense A.P.F.R.L. (2009). Frequency of tumors in dogs based on cytological diagnosis: a retrospective study in a veterinary teaching hospital. *Semina: Ciências Agrárias, Londrina.* 30: 189-200.
- Šoštarić-Zuckermann I.C., Severin K., Hohšteter M., Artuković B., Beck A., Gudan Kurilj A., Sabočanec R., Džaja P., Grabarević Ž. (2013). Incidence and types of canine tumours in Croatia. *Vet. Arhiv.* 83: 31- 45.
- Tjalma R.A. (1966). Canine bone sarcoma. Estimation of relative risk as a function of body size. *J. Natl. Cancer Inst.* 36: 1137-1150.
- Vascellari M., Baioni E., Ru G., Carminato A., Mutinelli E. (2009). Animal tumour registry of two provinces in northern Italy: incidence of spontaneous tumors in dogs and cats. *BMC Vet. Res.* 5: 39.
- VetCancer Registry <http://www.vetcancerregistry.com> [accessed 10 April 2019].
- VMDB (2010). <https://vmdb.org/> [accessed 26 April 2019].