

Effect of proximity to Jujube (*Ziziphus lotus* and *Ziziphus jujube*) trees on medfly (*Ceratitis capitata*) populations in citrus orchards of Moulouya Perimeter

Jamal BEN YAZID

Zouheir CHAFIK

Imane BIBI

Anwar BOUSAMID

Ez-Zahra KHARMACH

Laboratory of Biochemistry and Biotechnology, Mohamed first University, Faculty of Sciences, Oujda, Morocco

Laboratory of Plants and Microorganisms Biology, Mohamed first University, Faculty of Sciences, Oujda, Morocco

Laboratory of Biochemistry and Biotechnologies Mohamed first University, Faculty of Sciences, Oujda, Morocco

Laboratory of Biochemistry and Biotechnologies Mohamed first University, Faculty of Sciences, Oujda, Morocco

Laboratory of Biochemistry and Biotechnologies Mohamed first University, Faculty of Sciences, Oujda, Morocco

Moulouya Perimeter is one of the most important zones of Citrus production in Morocco, with an area of 22.000 ha and 360.000 tons of Citrus production. Medfly "*Ceratitis capitata*, Wiedemann 1824" is a pest with an important economic impact for Citrus, because of its impact on yield losses and phytosanitary concern linked to its presence. In Citrus orchards of Moulouya perimeter, adjacent wild jujube (*Ziziphus lotus* and *Ziziphus jujuba*) trees may have an impact on Medfly populations. The main objective of this study is to evaluate the influence of adjacent jujube trees, growing in windbreaks and noncommercial orchards, on medfly captures in the traps located in citrus orchards and to determine the amount of increase of punctured fruits caused by medfly oviposition considering the distance ranges to the jujube trees (10, 25, 50, 75, 100, 125, 150, 175, 200, 225, 250 and 275m). To achieve this purpose, 180 traps were placed in 15 orchards and monitored for medfly captures weekly, from April to November 2019, within three locations (Aklim, Chouihia, and Reggada) representing Citrus producing area of Moulouya Perimeter. The monitoring of *C. capitata* adults was conducted using "Maghreb med" traps with trimedlure (tbutyl-4(or5)-chloro-2-methyl-cyclohexane carboxylate) male-specific para-pheromone as attractant and dichlorvos as a toxicant. Five orchards per each location were selected and 12 traps were hung in every orchard in fixed distances from the jujube trees and fruits were checked for punctures. The influence of each jujube tree in citrus orchards in his vicinity is higher within distances inferior to 75 m. Then, its impact could approach 2,25 Ha on average, and in this area, *C. capitata* population densities were twice to five times as high as in other distances. Furthermore, Citrus trees located at distances inferior to 75m from the jujube trees showed an increase in the percentage of punctured fruits compared with distances higher than 75m.

INTRODUCTION

The Mediterranean fruit fly (hereafter medfly), *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae) is one of the most devastative insect pests of fruit and vegetables worldwide (Papadopoulos et al., 2001). Adult females of medfly lay eggs in ripe or ripening fruits and larvae feed by destroying the fruit mesocarp (pulp, flesh). Certain studies pointed out that the presence of jujube, patches or trees, in cropped lands constitutes a refuge for several crops pests (weeds, mollusk, rodents, Spanish sparrow, insects and bacteria) (Rsaissi et al., 2012). Non-cultivated,

isolated jujube trees in windbreaks and adjacent noncommercial orchards are usually present in large numbers in the landscape of citrus crops in the Moulouya perimeter, like in other Mediterranean areas (Roselló i Botey, 2007). Mature jujube fruits are frequently attacked by female medfly causing punctures and serving as multiplication media and origin of new infestations in other cultivated crops like Citrus (Ben yazid et al. 2018). Several studies reported that adjacent host plants to Citrus commercial orchards are a reservoir of infestation of those orchards (Israely et al., 1997; Alemany et al., 2004; Martínez-Ferrer et al., 2006). Because of the high risk of infestation, certain authors recommend considering the control of *C. capitata* in them (Muñoz and Cardeñoso Herrero, 2006). The main objective of this study is to check the influence of adjacent jujube trees on *C. capitata* captures and fruit damage in citrus orchards considering the ranges of distances between citrus trees and traps from the jujube trees.

MATERIAL AND METHODS

Study area

The study was carried out in three sites in the citrus producing area of Moulouya Perimeter, in northeast Morocco, from May to November 2019 (Figure 1). Five orchards per location (Aklim, Chouihia, and Reggada) commercial clementine (*Citrus clementina*) orchards, with adult trees, were selected for the survey (the criteria of selection were jujube trees presence in vicinity and the presence of clementine trees) (Figure 1). Selected orchards were representing the total Citrus growing area. The monitoring of *C. capitata* adults flies was based on “Maghreb med” traps with trimedlure (tbutyl-4(or5)-chloro-2-methyl-cyclohexane carboxylate) male-specific parapheromoun as attractant and dichlorvos (DDVP) as a toxicant agent. To maintain the effectiveness of traps, Trimedlure ships were changed every 8 weeks, and DDVP ships every 6 weeks. In each orchard, 12 traps were placed and set up at 1,5 m above the ground on the sunny side of the tree. The captures of *C. capitata* were checked in the 180 traps weekly and noted, captured flies were removed from the traps after counting and noting. To avoid interferences, traps were hung at 50 m away from ones each other and in orchards without the presence of other host plants than jujube in hedges.

The amounts of captures and fruit punctured were compared based on the distance to the jujube trees growing in windbreaks or noncommercial adjacent orchards. Traps were placed at fixed distances from the jujube trees. To ensure the measurement of distances between traps and the nearest jujube trees, all surveyed trees for punctured fruits, all the traps, and jujube trees in the windbreak or noncommercial orchards were georeferenced and marked. Retained distances were 10; 25; 50; 75; 100; 125; 150; 175; 200; 225; 250; 275 meters from the jujube trees.




Figure 1: Location of study area sites in Moulouya Perimeter North east of Morocco (Aklim, Chouihia and Reggada)

Data and statistical analysis

Captures and punctured fruits in different locations, orchards and traps were synthesized and compared using General Linear Model Univariate (GLM) and means were separated with Post hoc test Scheffé_{a,b} with $P \leq 0.05$ and 95% confidence intervals (CI). All data statistical analyses were carried out using the SPSS.25 software and Excel Microsoft.

RESULTAS AND DISCUSSION

Fluctuations of medfly populations in the study area

Medfly captures occurred, during our study period, since the second week of April and the peak was recorded earlier in September. After the peak, the number of captures started to decrease and attained zero at the end of November (Figure 2). The same results have been reported by Ben yazid et al. in 2020. This fluctuation is linked to climatic conditions and the availability of host plants and ripening fruits in the wild and cultivated fruit trees. The period between late August and early September is a key period for citrus fruits because it corresponds to the ripening stage which is with high sensitivity to medfly punctures (Ben yazid et al., 2018).




Figure 2: Medfly captures in FTD (Flies per trap and day), in all traps during the study period

Effect of proximity to jujube trees on medfly captures in Citrus orchards

The analysis of collected data showed that captures are influenced by the distance of traps from the jujube trees. In three locations (Reggada, Aklim, and chouihia), the pattern of distribution of captures in all traps presented a significant difference (Table 1). The means of medfly captures were 316,1; 371,75 and 423,73 with error standard of 17,106 for Aklim, Chouihia, and Reggada respectively. This result is due to the prevalence of jujube trees in Reggada location and the abundance of the host plant in non-controlled orchards.

Independent variable: Medfly Captures				
Location	Mean	Standard error	Confidence interval at 95 %	
			Lower bound	Upper bound
Aklim	316,100	17,106	282,321	349,879
Chouihia	371,750	17,106	337,971	405,529
Reggada	423,733	17,106	389,955	457,512

Table 1: Estimated averages of medfly captures in three study locations Aklim, Chouihia and Reggada

Our results (Figure 3) showed that the traps located within a distance of 75 m from a jujube tree captured the highest level of captures in all locations and all orchards. Those traps showed a double to five times increase in total number of flies captured per trap during study period, compared with traps located at other considered distances higher than 75m. This increase in captures was important. All the distance ranges considered between 75m < distances < 275m showed lower captures compared with traps located at ≤ 75 m from jujube trees in all orchards and locations. Results may explain that jujube trees are a reservoir of infestations of citrus orchards and that captures in Citrus orchards are significantly linked to the distance between traps and jujube trees present in windbreaks or adjacent noncommercial orchards. The Table 2 showed that traps in distances inferior or equal to 75m from the jujube trees are grouped in a three sub-sets

with means of 438,66bcd ; 515,73cd;656,33de; 777,46e flies per trap in 75; 50; 25 and 10 m of distance significantly different to the mean of other sub-sets (257.26 flies per trap) which grouping traps representing all other distances from jujube trees (75<distances<275 with some slight differences inter sub-sets) with means of varying within interval of 146 and 394,93 flies per trap.

Considering the influence of each jujube tree in citrus orchards in his vicinity is higher within distances inferior to 75 m, this impact approaches 2,25 Ha on average and in this area *C. capitata* population density could be twice to five times as high as in other areas. In Molouya perimeter, the chemical spray decision making, for Medfly control, is piloted only by the threshold of male adults captures in traps hung on citrus trees in orchards. However, medfly captures are linked to the presence of host fruit trees in windbreaks and adjacent hedges surrounding the citrus orchards (mainly jujube trees). Thus, the presence of jujube trees in our study zone and noncommercial orchards surrounding citrus orchards provides a potential breeding area for generations that will later attack Clementine varieties in later September, October, and earlier November. To reduce this effect, it is recommended to remove the host plants adjacent to citrus orchard, integrate them in the control programs. Furthermore its might be useful as “trap tree” for medfly trapping. Our results concord with those obtained for fig trees in Ibiza Island in Spain by Muñoz et al. in 2011 with an average of the impact of 1 Ha and distance of trap from fig tree of 50 m. Martínez-Ferrer et al. (2006) reported the same influence for other host plants adjacent to citrus orchards like peach and apricot trees in Spain.




Figure 3: Influence of distances of traps from jujube trees on total medfly captures per trap during the period of survey in the three locations (Aklim, Chouihia and Reggada).

Post hoc test Scheffé ^{a,b}		
Distance of Traps from jujube tree	Number of traps	Means (flies per Trap)
275m	15	146,00 ^a
250m	15	166,46 ^a
225m	15	205,26 ^{ab}
200m	15	230,33 ^{ab}
175m	15	266,93 ^{abc}
150m	15	300,86 ^{abc}
125m	15	347,33 ^{abc}
100m	15	394,93 ^{abc}
75m	15	438,66 ^{bcd}
50m	15	515,73 ^{cd}
25m	15	656,33 ^{de}
10m	15	777,46 ^e

Table 2: Mean of total captures in all traps with all studied distances (Values with the same letter do not differ GLM and Scheffé a,b test ($P < 0,05$; $F=33,303$; $P \text{ value} = ,000$; $df=(11; 162)$) Calculated Means are based on observed means; Error (Mean square) = 24108,155; $N= 15$; $\alpha=0,05$ and Significances were: 0,066a; 0,121b; 0,67c; 0,207d; 0,948e)

Effect of proximity to jujube trees on punctured fruits caused by medfly in Citrus orchards

The percentage of affected Citrus fruits was under 5% in the trees under survey because of intensive management of medfly in the area by the growers. But this percentage still presents high risk of interception of larvae in exported shipments. Data showed that the mean percentage of affected Citrus fruits was under 5% and the rate of punctured fruits was influenced by the distance of the citrus tree surveyed from the jujube trees. In the three locations, the pattern of distribution of punctures in all surveyed trees showed a significant difference between Reggada and Aklim, and between Reggada and Chouihia, but a significant similarity between Aklim and Chouihia (Figure 4). The distribution within each location and orchard is influenced by the distance of the citrus tree from the jujube tree. The results of our study showed an increase in punctured fruit as Citrus trees were close to the jujube trees. Subsets with less than 75m distance from jujube trees showed higher infestation rates compared to subsets with distances longer than 75m with a high significance (Table 3). The mean percentage of punctured fruits, in subsets within distances lower than 75m, was 4 to 8 times higher than mean percentage in other distances.

The fruit punctures trends found in all locations in citrus zone were similar to those reported for other Mediterranean countries or regions, with only relatively minor variations (Muñoz et al., 2011).




Figure 4: The effect of distances of traps from jujube trees on percentage of fruit punctured in surveyed Citrus trees in the three locations (Aklim, Chouihia and Reggada)

Post hoc test Scheffé ^{a,b}		
Distances of citrus trees from jujube trees	Number of trees	Means (Percentage Of punctured fruits %)
275m	15	0,83 ^a
250m	15	0,93 ^{ab}
225m	15	1,13 ^{ab}
200m	15	1,53 ^{abc}
175m	15	1,66 ^{abc}
150m	15	1,90 ^{bcd}
125m	15	2,36 ^{cde}
100m	15	2,43 ^{cde}
75m	15	2,80 ^{def}
50m	15	3,33 ^{efg}
25m	15	3,73 ^{fg}
10m	15	4,13 ^g

Table 3: Mean of percentages of punctured citrus fruits in all studied distances (Values with the same letter do not differ referring to GLM and Scheffé^{a,b} test ($\alpha = 0,05$; $F=11,644$; $P \text{ value} = 0,000$; $df = (4; 162)$). Calculated means are based on observed means. Error (Mean square) = 0,345; $N = 15$; $\alpha = 0,05$ and Significances were 0,191a; 0,050b; 0,103c; 0,103d; 0,050e; 0,073f; 0,249g)

CONCLUSION

Our findings suggest that adjacent non-cultivated jujube trees growing in windbreaks and noncommercial orchards reinforced the increase of medfly captures and contributed to increasing the damage in citrus fruits adjacent to those isolated plants. This result stresses the convenience of including jujube trees in medfly populations control programs, when it is implemented, in citrus orchards located in their vicinity. Also, Jujube trees might be useful as "trap trees" in medfly control programs if they can't be removed (Martínez-Ferrer et al., 2006). Extending trials are recommended to other host plants present in the vicinity of citrus orchards, mainly pomegranate, loquat, fig, and prickly pear trees.

REFERENCES

- Aleman A., Miranda M.A., Alonso R., Martín Escorza C. (2004). Efficacy of *C. capitata* (Wied.) (Diptera: Tephritidae) female mass trapping. Edge-effect and pest multiplier role of unmanaged fruit hosts. *Bol. San. Veg. Plagas*, 30: 255-264.
- Ben yazid J., Chafik Z., Bousamid A., Bibi I., Kharmach E. (2020). Population Dynamics and Seasonal Occurrence of Mediterranean Fruit Fly (*Ceratitis capitata* Wiedemann, 1824) in Moulouya Perimeter North East of Morocco. *Indian Journal of Ecology* 47(2): 564-569.
- Ben yazid J., Chafik Z., Bousamid A., Bibi I., Kharmach E. (2018) Current situation and seasonal occurrence of Mediterranean fruit fly (*Ceratitis capitata*, Wiedmann, 1824) in Moulouya Perimeter North east of Morocco, in: Onzième Congrès de l'Association Marocaine de Protection des plantes 26-27 Mars 2018, Rabat, Maroc: Proceeding : 213-219.
- Israely N., Yuval B., Kitron U., Nestel D. (1997). Population fluctuations of adult Mediterranean Fruit Flies (Diptera: Tephritidae) in a mediterranean heterogeneous agricultural region. *Environmental Entomology*, 26(6): 1263-1269.
- Martínez-Ferrer M.T., Campos Rivela J.M., Fibla Queral J.M. (2006). Population dynamics of *Ceratitis capitata* on Citrus in northeastern Spain: influence of adjacent host fruit trees. *IOBC/WPRS Bull*, 29(3): 77-84.
- Muñoz A.A., Cardeñoso Herrero M. (2006). La mosca del Mediterráneo. *Món Rural d'Eivissa i Formentera. La revista de la Fira del camp*, 4: 20-21.
- Muñoz A.A., Garcia-Marí F. (2011). Proximity to fig trees increases medfly populations in citrus orchards. *IOBC/wprs Bulletin*, 62: 229-233.
- Papadopoulos N.T., Katsoyannos B.I., Carey J.R., Kouloussis N.A. (2001). Seasonal and annual occurrence of the Mediterranean fruit fly (Diptera: Tephritidae) in northern Greece. *Annals of the Entomological Society of America*, 94(1): 41-50.
- Rosselló i Botey J. (2007). Caracterizació de cultivars de Figuera a Mallorca. Govern de les Illes Balears. Conselleria d'Agricultura i Pesca. Quaderns d'Agricultura, 12: 175.
- Rsaissi N., Bouhache M., Bencharki B. (2012). Importance and agro-economic impact of wild jujube (*Ziziphus lotus*) in Chaouia region. *Revue Marocaine de Protection des Plantes*, 3 : 13-27.

References