

First report of *Drosophila suzukii* (Diptera: Drosophilidae) in North Africa

M'bark OUARTAR

National School of Applied Sciences (ENSA), Ibn Zohr University, Agadir, Morocco

Gianfranco ANFORA

Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige (TN), Italy

Rachid BOUHAROU

Regional Center of Agronomic Research (INRA), Agadir, Morocco

Bouchra CHEBLI

National School of Applied Sciences (ENSA), Ibn Zohr University, Agadir, Morocco

Spotted wing *Drosophila suzukii* (Diptera: Drosophilidae) is an important invasive pest reported in the last few years in Europe. The female lays eggs preferably on soft fruits like cherries and others causing thereafter injury by development of rotting point. Red traps baited with apple cider vinegar, red wine and sugar were used to collect the *D. suzukii* adults. After preparation of slide mounting, the key identification was used to confirm presence of this pest in the north of Morocco. The dark spotted wings of male and sclerotized black and strong teeth on valve margin of female ovipositor were observed. Until early 2014, damages were not detected but the reviewing and updating of pest risk assessment should be implemented to prevent heavy economic losses.

Keywords: *Drosophila suzukii*, soft fruits, pest risk assessment

Introduction

Drosophila suzukii known as the Spotted Wing *Drosophila* (SWD) (Matsumura) (Diptera: Drosophilidae) is a new invasive pest native of South East Asia. In 2008, the SWD was detected simultaneously in North America and Europe, where this pest caused a serious damage to fruit production the following few years (Cini, 2012; Lee et al., 2011; Pfeiffer et al., 2012; Walsh et al., 2011).

Currently, the vast majority of *Drosophila* species, commonly known as vinegar flies, has become cosmopolitan species after colonizing new areas in the globe (Fernandez Iriarte et al., 2009) although most of them are not considered harmful pests as their larvae mostly develop in injured and rotting fruits. In contrast to these species, SWD exhibits a wide variety of host crops and ripening wild and cultivated fruits. Shortly before harvest, *Drosophila suzukii* lays its eggs in ripening fruits such as cherries, strawberries, blueberries, raspberries, blackberries, apricots, peaches, grapes and a number of wild fruits. Therefore, healthy unwounded fruits are favored to overripe ones (Mitsui et al., 2006). The *Drosophila suzukii* hence causes important economic losses (Lee et al., 2011; Walsh et al., 2011; De Ros et al., 2013). Females are exceptionally fecund (400 eggs/female), the life cycle from egg hatching to adult emergence ranges from about 9-10 days to 21-25 days respectively at 25° and 15°C, and hence from 7 to 13 generations can carry out per year (Kanzawa, 1939).

Drosophila suzukii adults are brown-yellow small (<4 mm) flies with red eyes. The abdominal segments possess unbroken dark stripes and the last segment is often dark. Males and females can be distinguished by their sexual dimorphism (Pfeiffer et al., 2012). Males possess a single dark spot on the leading edge of each wing. On the first and second segments of their front legs, males also exhibit two black combs with 3 to 6 teeth parallel to the length of the legs. In contrast, females are

deprived of these characteristics but possess a large serrated ovipositor that they use to lacerate the fruit skin. Eggs, larvae and pupae are difficult to identify due to their similarities with other species (Cini et al., 2012).

Material and Methods

On June, 2013, a single male of the SWD, was captured accidentally in yellow sticky traps in Moulay Bouselham, Kenitra Province (Morocco).

Subsequently in October, red traps were placed in a raspberry orchard; each trap consists of a plastic container, baited with 150 ml of apple cider vinegar, 50 ml of red wine and one spoon of sugar, with small holes of 4 mm diameter allowing the entrance of the attracted flies (Grassi and Maistri, 2013). Twenty adults of SWD (male and female) were collected and subjected to identification using dichotomous key for identification of *Drosophila suzukii* (Hauser, 2011; Vlach, 2012) at the plant protection laboratory of Regional Center of INRA (Institut National de la Recherche Agronomique) in Agadir-Morocco and a voucher specimen is taken in the same laboratory.

Results and discussion

The SWD male can be recognized based on the spots in their wing (Figure 1). The female possesses a long and narrow ovipositor with sclerotized black teeth (Figure 2 and Figure 3).



Figure 1: General view of *Drosophila suzukii* male. The black spots can be observed in the male's wings

Figure 1:



Figure 2: ovipositor of Spotted Wing Drosophila

Figure 2:



Figure 2: ovipositor of Spotted Wing Drosophila

Figure 3:

The first reported detection of *D. suzukii* in the North of Mediterranean basin was in 2008 in Rasquera, Spain (Calabria et al., 2012). Until early 2013, Mediterranean detection of SWD had been limited to Europe. The discovery in Morocco is of great importance since these are the first records of the species in the more dry Mediterranean climates. Ecological simulations seem to indicate that those Mediterranean conditions are not optimal for the growth of *D. suzukii* (Mitsui et al., 2006).

SWD may be introduced to Moroccan fields as adult flies associated with fruit, or as eggs, larvae or pupae within fruit. Indeed, the proximity of Morocco to Spanish coast (about 14 km) and the occurring of an intensive international trade between Morocco and Europe could be a rationale.

Unlike most other members of the Drosophilidae family, this highly fecund species (7 to 13 eggs per day, Kanzawa 1939) is able to infect healthy ripening fruit instead of overripe and rotting fruit. *D. suzukii* can lay eggs on healthy soft bodied fruit and can even attack hard fruit crops such as peaches, apples and pears (Walsh et al., 2011). After infestation, secondary parasites and invaders often contribute to fruit deterioration. These facts make SWD an economically serious pest that needs management at highest level possible.



Because this pest is widespread in the world as it occurs in North America, Europe and Asia, it is considered non-actionable; meaning international trade restrictions or quarantines will be implemented.

Trapping is highly required for the early detection and management of *D. suzukii*. Mass trapping is less costly tool and easy to perform compared to other methods of control. An early detection of this potential pest is decisive for good management of host crops and monitoring effort is necessary to control the probable further expansion and/or establishment of this species in new areas in the country. The local quarantine measures are worth considering. Until early 2014, damages were not detected but the reviewing and updating of pest risk assessment should be implemented to prevent heavy economic losses.

References

- Walsh D.B., Bolda M.P., Goodhue R.E., Dreves A.J., Lee J., Bruck D.J., Walton V.M., O'Neal S.D., Zalom F.G. (2011). *Drosophila suzukii* (Diptera:Drosophilidae): invasive pest of ripening soft fruit expanding its geographic range and damage potential. *J. Integrated Pest Manag.*, 1: 1-7.
- Calabria G., Máca J., Bächli G., Serra L., Pascual M. (2012). First records of the potential pest species *Drosophila suzukii* (Diptera: Drosophilidae) in Europe. *J. App. Entomol.*, 136: 139-147.
- Cini A., Ioriatti C., Anfora G. (2012). A review of the invasion of *Drosophila suzukii* in Europe and a draft research agenda for integrated pest management. *Bull. Insectol.*, 65: 149-160.
- De Ros G, Anfora G, Grassi A, Ioriatti C (2013) The potential economic impact of *Drosophila suzukii* on small fruits production in Trentino (Italy). *IOBC/WPRS Bull.*, 91: 317-321.
- Fernandez Iriarte P., Balanyà J., Pascual M., Mestres F., Hasson E., Fontdevila A., Serra L. (2009). Tracking the origin of an invasive species: *Drosophila subobscura* in Argentina. *J. Evolut. Biol.*, 22: 650-658.
- Grassi A., Maistri S. (2013). *Drosophila suzukii* su piccoli frutti e ciliegio. *Terra Trentina*, 3: 47-53.
- Hauser M. (2011). A historic account of the invasion of *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) in the continental United States, with remarks on their identification. *Pest Manag. Sci.*, 67: 1352-1357.
- Kanzawa T. (1939). Studies on *Drosophila suzukii* Mats. Kofu, Yamanashi agricultural experiment station 49 pp. Abstract in *Rev. Appl. Entomol.*, 29: 622.
- Lee J.C., Bruck D.J., Curry H., Edwards D., Haviland D.R., Van Steenwyk R.A., Yorgey B.M. (2011). The susceptibility of small fruits and cherries to the spotted-wing drosophila, *Drosophila suzukii*. *Pest Manag. Sci.*, 67: 1358-1367.
- Mitsui H., Takahashi K.H., Kimura M.T. (2006). Spatial distributions and clutch sizes of *Drosophila* species ovipositing on cherry fruits of different stages. *Popul. Ecol.*, 48: 233-237.
- Pfeiffer D.G., Leskey T.C., Burrack H.J. (2012). Threatening the harvest: The threat from three invasive insects in late season vineyards. In: Bostanian N.J., Vincent C., Isaacs R. (eds), *Arthropod Management in Vineyards: Pests, Approaches, and Future Directions*. Springer, Dordrecht, The Netherlands, pp 449-474.
- Vlach J. (2012). http://www.agf.gov.bc.ca/cropprot/swd_identification.pdf Accessed 28 June 2013.



References